arXiv:1211.7087v1 [math.AT] 29 Nov 2012

When does simplicial homology vanish?

E. Connon[*]

December 3, 2012

Department of Mathematics and Statistics, Dalhousie University, Canada

Abstract

In this paper we examine the relationship between the combinatorial structure of a simplicial complex and its simplicial homology. In particular we introduce the combinatorial notion of a d-dimensional cycle and show that over any field of characteristic 2 the existence of non-zero d-dimensional homology corresponds exactly to the presence of a d-dimensional cycle in the simplicial complex. We also show that d-dimensional cycles give rise to non-zero simplicical homology over any field.

Keywords: simplicial homology, support complex, d-dimensional cycle, graph cycle, Stanley-Reisner ideal, pseudo d-manifold

1 Introduction

Simplicial homology is the oldest and most explicit of the homology theories. In algebraic topology it is used as a building block in the development of singular homology, but its influence extends into commutative algebra and combinatorics where it has deep implications in resolutions of modules and other algebraic properties. In [5] Hochster gives a fundamental formula for the betti numbers of the Stanley-Reisner ideal of a simplicial complex in terms of the dimensions of certain homology groups of the complex. A theorem of Reisner from [8] gives a condition for the Stanley-Reisner ring of a simplicial complex to be Cohen-Macaulay in terms of the simplicial homology of the complex and some of its subcomplexes. This theorem is used by Stanley in [9] to prove the upper bound theorem for simplicial spheres.

Despite the substantial use of simplicial homology in classifying algebraic properties and the concrete combinatorial nature of the theory there appears to be very little literature on the explicit combinatorial structures necessary for a simplicial complex to exhibit non-zero simplicial homology. However studies have been made into the combinatorics of acyclic simplicial complexes – those for which simplicial homology vanishes in all dimensions. In [6] Kalai gave a characterization of the f-vectors of such simplicial complexes. This was followed by work of Stanley in [10] on a combinatorial decomposition of these complexes. On the other hand the literature

---

[*] Research supported by NSERC and Killam scholarships.

relating to the combinatorial structures associated with non-zero simplicial homology is scant. In [3] Fogelsanger studied the rigidity of the 1-skeletons of minimal $(d-1)$-cycle complexes and concluded that these 1-skeletons are rigid graphs when embedded in $\mathbb{R}^d$. However the question remains as to whether or not one may attribute the existence of non-zero homology in a simplicial complex to a combinatorial structure present in that complex. It is this problem which we address in this paper. This is an interesting mathematical question in its own right, but at the same time its possible solution has the potential to enable easier translation between algebraic properties of monomial ideals and the combinatorial framework. For example, in [2] the results of this paper are used to give an explicit combinatorial condition that is satisfied whenever a Stanley-Reisner ideal has a linear resolution.

The author is grateful for comments of Manoj Kummini.

2 Preliminaries

2.1 Simplicial complexes and simplicial homology

An (abstract) **simplicial complex** Γ on the finite **vertex set** V is a set of subsets of V such that $\{v\} \in \Gamma$ for all $v \in V$ and for any $F \in \Gamma$ if $G \subseteq F$ then $G \in \Gamma$. The elements of V are **vertices** of Γ and the elements of Γ are called **faces** or **simplices** of Γ. Faces of Γ that are maximal with respect to inclusion are called **facets** of Γ.

The dimension of a face F of Γ, denoted $\dim F$, is defined to be $|F|-1$ while the dimension of Γ itself is given by

$$\dim \Gamma = \max\{\dim F : F \in \Gamma\}.$$

A face of dimension n is called an n**-face** or an n**-simplex**. If all facets of Γ have the same dimension then Γ is said to be **pure**.

To any d-face in a simplicial complex we can assign an **orientation** by specifying an ordering of its vertices. Two orientations are said to be **equivalent** if one is an even permutation of the other. Thus there are only two equivalence classes of orientations for each face. By an **oriented** d**-face** we mean a d-face with a choice of one of these orientations. We denote the d-face on vertices $v_0, \dots, v_d$ with the orientation $v_0 < \cdots < v_d$ by $[v_0, \dots, v_d]$.

Let A be a commutative ring with unit. Then we define $C_d(\Gamma)$ to be the free A-module whose basis is the oriented d-faces of Γ and where $[v_0, v_1, \dots, v_d] = -[v_1, v_0, \dots, v_d]$. The elements of $C_d(\Gamma)$ are called d**-chains**. The **support complex** of a d-chain is the complex whose facets are the d-faces in the d-chain.

There is a natural boundary map homomorphism ∂_d from the space of d-chains to the space of $(d-1)$-chains defined by setting

$$\partial_d([v_0, \dots, v_d]) = \sum_{i=0}^{d} (-1)^i [v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_d]$$

for each oriented d-face $[v_0, \dots, v_d]$. The kernel of ∂_d is called the group of d**-cycles** and the image of ∂_d is called the group of $(d-1)$**-boundaries**. The dth simplicial **homology group** of Γ over A is equal to the quotient of the group of d-cycles over the group of d-boundaries and is denoted $H_d(\Gamma; A)$. In essence a non-zero element of $H_d(\Gamma; A)$ corresponds to a “d-dimensional hole" in Γ. For example, when Γ is a triangulation of the sphere, which has a 2-dimensional hole, $H_2(\Gamma;\mathbb{Z})$ is non-zero.

We can also define a surjective homomorphism $\epsilon : C_0(\Gamma) \to \mathbb{Z}$ by $\epsilon(v) = 1$ for each $v \in V(\Gamma)$. The **reduced homology group** of Γ in dimension 0, denoted $\tilde{H}_0(\Gamma;A)$, is defined by

$$\tilde{H}_0(\Gamma;A) = \ker \epsilon / \text{im } \partial_1.$$

We set $\tilde{H}_i(\Gamma;A) = H_i(\Gamma;A)$ for $i > 0$ to obtain the ith **reduced homology group** of Γ. For a more in-depth explanation of simplicial homology see [7].

2.2 Graph Theory

A **finite simple graph** G consists of a finite set of **vertices** denoted $V(G)$ and a set of **edges** denoted $E(G)$ where the elements of $E(G)$ are unordered pairs from $V(G)$. In this paper we will use the term **graph** to refer to a finite simple graph. A 1-dimensional simplicial complex can be thought of as a graph where the edges are given by the 1-faces of the simplicial complex.

A vertex v of a graph G is said to be **incident** with an edge e of G if v is a vertex of e. The **degree** of a vertex v is the number of edges with which v is incident.

A **path** in a graph G is a sequence of vertices $v_0, \ldots, v_n$ from $V(G)$ such that $\{v_{i-1}, v_i\} \in E(G)$ for $1 \leq i \leq n$. A **connected** graph is a graph in which every pair of vertices is connected by a path.

A **cycle** in a graph G is an ordered list of distinct vertices $v_0, \ldots, v_n$ from $V(G)$ where $\{v_{i-1}, v_i\} \in E(G)$ for $1 \leq i \leq n$ and $\{v_n, v_1\} \in E(G)$. In Figure 1 we give an example of a graph cycle on six vertices.



Figure 1: Graph cycle on six vertices.

3 Motivation

Many of the homological classifications used in commutative algebra are based on the vanishing of simplicial homology in certain simplicial complexes. We would like to turn this property into an explicitly combinatorial one and so our goal is to relate the concept of non-zero homology in a simplicial complex to the existence of a specific combinatorial structure in that complex which is distinct from the more algebraic notion of a d-chain.

It is not difficult to show that this goal is achievable for non-zero simplicial homology in dimension 1. In this case, the combinatorial structure associated to non-zero homology is the graph cycle. This can be deduced from [1, Chapters 4 and 5], but in this section we provide an explicit proof of this relationship as motivation for the general case. The techniques used in this proof also illustrate our overall approach to this problem.

First we recall the following well-known lemma from graph theory. A proof can be found in [11, Section 1.2].

Lemma 3.1. *If G is a finite simple graph in which every vertex has degree at least 2 then G contains a cycle.*

Theorem 3.2. *For any simplicial complex Γ and any field k, $\tilde{H}_1(\Gamma; k) \neq 0$ if and only if Γ contains a graph cycle, which is not the support complex of a 1-boundary.*

Proof. Suppose that $\tilde{H}_1(\Gamma; k) \neq 0$. Then Γ contains a 1-cycle c that is not a 1-boundary. We may assume that the support complex of c is minimal with respect to this property. In other words no strict subset of the 1-faces of c is the support complex of a 1-cycle which is not a 1-boundary. First we would like to show that the support complex Ω of c is a connected graph.

The 1-cycle c is of the form

$$c = \alpha_1 F_1 + \cdots + \alpha_n F_n \tag{1}$$

for some oriented 1-faces $F_1, \ldots, F_n$ of Γ and where $\alpha_i \in k$. If Ω is not connected then we can partition the 1-faces $F_1, \ldots, F_n$ into two sets having no vertices in common. Without loss of generality let these two sets be $\{F_1, \ldots, F_\ell\}$ and $\{F_{\ell+1}, \ldots, F_n\}$. Since

$$\partial_1(c) = \partial_1(\alpha_1 F_1 + \cdots + \alpha_n F_n) = 0$$

and since there are no vertices shared between the two sets we must have

$$\partial_1(\alpha_1 F_1 + \cdots + \alpha_\ell F_\ell) = 0 \text{ and } \partial_1(\alpha_{\ell+1} F_{\ell+1} + \cdots + \alpha_n F_n) = 0$$

and so $\alpha_1 F_1 + \cdots + \alpha_\ell F_\ell$ and $\alpha_{\ell+1} F_{\ell+1} + \cdots + \alpha_n F_n$ are both 1-cycles. By the assumption of minimality of the support complex of c we know that these 1-cycles must also be 1-boundaries. However, since c is the sum of these two 1-chains and they are both 1-boundaries c must be a 1-boundary as well, which is a contradiction. Therefore Ω must be a connected graph.

Next, note that the degree of all vertices in Ω must be at least two. This follows since $\partial_1(c) = 0$ and this may only be achieved if all vertices present cancel out in this sum. Therefore each vertex must appear at least twice. Hence by Lemma 3.1 we know that Ω contains a graph cycle. Let $v_0, \ldots, v_m$ be the vertices in this cycle where v_i is adjacent to v_{i+1} for $0 \leq i \leq m-1$ and v_m is adjacent to v_0. By relabeling if necessary we may assume that $F_1, \ldots, F_m$ are the oriented 1-faces corresponding to the edges in this cycle with $F_i = \varepsilon_i[v_i, v_{i+1}]$ for $0 \leq i \leq m-1$ and $F_m = \varepsilon_m[v_m, v_0]$ where $\varepsilon_i = \pm 1$ depending on the orientation of F_i. Suppose that $m < n$. Then

$$\varepsilon_1 F_1 + \ldots + \varepsilon_m F_m = \sum_{i=0}^{m-1} [v_i, v_{i+1}] + [v_m, v_0]$$

is a 1-chain and it is straightforward to see that

$$\partial_1(\varepsilon_1 F_1 + \ldots + \varepsilon_m F_m) = 0.$$

Let $b = \alpha_1\varepsilon_1(\varepsilon_1 F_1 + \cdots + \varepsilon_m F_m)$. Then

$$\partial_1(b) = \alpha_1\varepsilon_1\partial_1(\varepsilon_1 F_1 + \ldots + \varepsilon_m F_m) = 0$$

and so b is a 1-cycle. Now $m < n$ and so by our assumption of minimality of the support complex of c we know that b is also a 1-boundary. We have

$$\partial_1(c - b) = \partial_1(c) - \partial_1(b) = 0 - 0 = 0$$

and so $c - b$ is a 1-cycle. Since

$$c-b = 0+(\alpha_2-\alpha_1\varepsilon_1\varepsilon_2)F_2+(\alpha_3-\alpha_1\varepsilon_1\varepsilon_3)F_3+\cdots(\alpha_m-\alpha_1\varepsilon_1\varepsilon_m)F_m+\alpha_{m+1}F_{m+1}+\cdots+\alpha_n F_n,$$

it is supported on a strict subset of the 1-faces of c and so by the assumption of minimality $c - b$ is a 1-boundary. Therefore, since both b and $c - b$ are 1-boundaries and $c = b + (c - b)$ then c is a 1-boundary also. This is a contradiction and so we must have $m = n$. Therefore Ω itself is a graph cycle.

Next we will show that Ω is not the support complex of a 1-boundary. To begin we show that for any 1-cycle of the form

$$d = \beta_1 F_1 + \cdots + \beta_n F_n$$

for non-zero $\beta_1, \ldots, \beta_n \in k$ we have $\beta_1 = \cdots = \beta_n$. Let v_1^i, v_2^i be the two vertices belonging to the 1-face F_i. Since d is a 1-cycle we have

$$0 = \partial_1(d) = \beta_1 v_1^1 - \beta_1 v_2^1 + \cdots + \beta_n v_1^n - \beta_n v_2^n \tag{2}$$

and since Ω is a graph cycle each v_s^t appears exactly twice in (2) as it belongs to two 1-faces. Therefore if F_i and F_j are any two 1-faces which share a vertex then we get $\beta_i = \beta_j$ from (2) and so any two 1-faces sharing a vertex have the same coefficient. Since Ω is connected we have $\beta_1 = \cdots = \beta_n$.

Therefore $\alpha_1 = \cdots = \alpha_n$ in (1) and if there exists a 1-cycle in Γ with Ω as a support complex which is also a 1-boundary then this implies that c is also a 1-boundary after multiplication by a constant in k. This is a contradiction and so Ω is a graph cycle which is not the support complex of a 1-boundary.

Conversely, suppose that Γ contains a graph cycle Ω, which is not the support complex of a 1-boundary. Let $v_0, \ldots, v_k$ be the vertices of Ω where v_i is adjacent to v_{i+1} for $0 \le i \le k - 1$ and v_k is adjacent to v_0. Then

$$c = \sum_{i=0}^{k-1}[v_i, v_{i+1}] + [v_k, v_0]$$

is a 1-chain whose support complex is Ω and it is easy to see that $\partial_1(c) = 0$. Therefore c is a 1-cycle, and by assumption it is not a 1-boundary. Therefore $\tilde{H}_1(\Gamma; k) \neq 0$.

□

4 d-dimensional cycles

We would like to be able to capture the idea of non-zero higher-dimensional simplicial homology by extending the idea of a graph cycle to higher dimensions. We first need the following definitions.

Definition 4.1 (**d-path, d-path-connected, d-path-connected components).** *A sequence $F_1, \dots, F_k$ of d-dimensional faces in a simplicial complex is called a d-***path** *between F_1 and F_k if for all $1 \leq i \leq k-1$ we have that $|F_i \cap F_{i+1}| = d$. A set of d-faces is d-***path-connected** *or* **strongly connected** *if there exists a d-path between each pair of its d-faces. The d-***path-connected components** *of a pure d-dimensional simplicial complex are the maximal subcomplexes which are d-path-connected.*

In Figure 2a we give an example of a 2-path between the 2-faces F_1 and F_2. Figure 2b shows a pure 2-dimensional simplicial complex with two 2-path-connected components shown by different levels of shading.



(a) A 2-path between F_1 and F_2.

(b) 2-path-connected components.

Figure 2

Definition 4.2 (**d-dimensional cycle).** *A pure d-dimensional simplicial complex Ω is called a d-***dimensional cycle** *if*

1. *Ω is d-path-connected, and*
2. *every $(d-1)$-face of Ω is contained in an even number of d-faces of Ω.*

The notion of a d-dimensional cycle extends the classical concept of a pseudo d-manifold that appears in algebraic topology. See for example [7].

Definition 4.3 (pseudo d-manifold). *A* **pseudo d-manifold** *Γ is a pure, d-dimensional d-path-connected simplicial complex in which every $(d-1)$-face of Γ is contained in exactly two d-faces of Γ.*

Examples of 2-dimensional cycles are given in Figure 3. Notice that the 2-dimensional cycles in Figures 3a through 3d are all examples of pseudo 2-manifolds while Figures 3e and 3f are not as they have 1-faces belonging to more than two 2-faces.

Definition 4.4 (**face-minimal d-dimensional cycle).** *A d-dimensional cycle Ω is called* **face-minimal** *if there is no d-dimensional cycle on a strict subset of the d-faces of Ω.*

Notice that a 1-dimensional cycle is a graph cycle if and only if it is face-minimal. In Figure 3 the only 2-dimensional cycle which is not face-minimal is 3e.

Lemma 4.5 (**A d-dimensional cycle can be broken into face-minimal ones).** *Any d-dimensional cycle Ω can be partitioned into face-minimal d-dimensional cycles $\Phi_1, \dots, \Phi_n$. In other words every d-face of Ω belongs to some Φ_i and no two distinct cycles Φ_i and Φ_j share a d-face.*



(a) A hollow octahedron.

(b) A triangulation of the real projective plane.

(c) A triangulation of the sphere.

(d) A triangulation of the torus.



(e) Two triangulated square pyramids glued along a 1-face.

(f) A triangulation of the sphere pinched along a 1-face.

Figure 3: Examples of 2-dimensional cycles.

Proof. If Ω is a face-minimal d-dimensional cycle then we are done. So suppose that Ω is not face-minimal and let Φ_1 be a face-minimal d-dimensional cycle on a strict subset of the d-faces of Ω. Consider the d-path-connected components of the complex Ω_1 whose facets are the d-faces of Ω not belonging to Φ_1. We claim that each such component is a d-dimensional cycle. Since each component is d-path-connected by definition we need only show that each $(d-1)$-face is contained in an even number of d-faces.

Let Ψ be one of the d-path-connected components of Ω_1 and let f be a $(d-1)$-face of Ψ. Suppose first that f also belongs to one of the d-faces of Φ_1. Since Φ_1 is a d-dimensional cycle f belongs to an even number of its d-faces. However Ω is also a d-dimensional cycle and f belongs to an even number of its d-faces. Since Ω_1 is the complex whose facets are the d-faces of Ω not in Φ_1 f belongs to an even number of d-faces in Ω_1. However the collection of all d-faces of Ω_1 containing f is clearly d-path-connected and so these d-faces all lie in Ψ. Hence f belongs to an even number of d-faces of Ψ.

If f does not belong to any d-faces of Φ_1 then all of the d-faces of Ω which contain f lie in Ω_1. Since there are an even number of such faces and they are d-path-connected they all lie in Ψ. Hence f belongs to an even number of d-faces of Ψ. Therefore each d-path-connected component of Ω_1 is a d-dimensional cycle.

Each of these components is either a face-minimal d-dimensional cycle, or contains a face-minimal d-dimensional cycle on a strict subset of its d-faces. We may repeat the argument above on the simplicial complex whose facets are the d-faces of Ω_1 from the components that are not face-minimal cycles. Iterating this procedure we see that since we have a finite number of d-

faces eventually the procedure must terminate. We are left with face-minimal cycles $\Phi_1, \ldots, \Phi_n$ in which every d-face of Ω belongs to some Φ_i and, by our construction, no two distinct cycles Φ_i and Φ_j share a d-face. ☐

Proposition 4.6. *A pseudo d-manifold is a face-minimal d-dimensional cycle.*

Proof. Let Ω be a pseudo d-manifold. Then Ω is d-path connected and every $(d-1)$-face in Ω belongs to exactly two d-faces. Hence Ω is a d-dimensional cycle. Suppose that Ω is not face-minimal. By Lemma 4.5 we can partition Ω into face-minimal d-dimensional cycles $\Phi_1, \ldots, \Phi_n$ where $n \geq 2$. Since Ω is d-path connected there must exist some pair of indices i, j with $i \neq j$ with $F_1 \in \Phi_i$ and $F_2 \in \Phi_j$ where $F_1 \cap F_2 = f$ for some $(d-1)$-face f of Ω. Since Φ_i is a d-dimensional cycle then f belongs to an even number of d-faces of Φ_i and similarly f belongs to an even number of d-faces of Φ_j. By Lemma 4.5, these d-faces are all distinct which means that f belongs to at least four d-faces of Ω. This is a contradiction since Ω is a pseudo d-manifold. Hence Ω is a face-minimal d-dimensional cycle. ☐

The converse of this theorem does not hold. As a counter-example see Figure 3f, a triangulated sphere pinched along a 1-dimensional face. It is a face-minimal 2-dimensional cycle, but it is not a pseudo 2-manifold as it has a 1-dimensional face belonging to four distinct 2-dimensional faces.

A pseudo d-manifold can be classified as either **orientable** or **non-orientable** and this idea can be generalized to the case of d-dimensional cycles. Recall from Section 2 that an orientation of a face in a simplicial complex is simply an ordering of its vertices.

Definition 4.7 (induced orientation). *Given an orientation of a d-face in a simplicial complex the* **induced orientation** *of any $(d-1)$-subface is given by the following procedure:*

- *if the vertex removed to obtain the $(d-1)$-face was in an odd position of the ordering then the orientation of the $(d-1)$-face is the same as the ordering of its vertices in the d-face*
- *if the vertex removed to obtain the $(d-1)$-face was in an even position of the ordering then the orientation of the $(d-1)$-face is given by any odd permutation of the ordering of the vertices in the d-face*

Definition 4.8 (orientable d-dimensional cycle). *We say that a d-dimensional cycle Ω with d-faces $F_1, \ldots, F_k$ is* **orientable** *if the following condition holds. There exists a choice of orientations of $F_1, \ldots, F_k$ such that for any $(d-1)$-face f of Ω when we consider the induced orientations of f by the F_i's containing f, these induced orientations are divided equally between the two orientation classes. Otherwise we say that Ω is* **non-orientable**.

Note that when we talk about the oriented d-faces of an orientable d-dimensional cycle we are referring to any set of orientations that is compatible with Definition 4.8.

The 2-dimensional cycles given in Figure 3 are all examples of orientable 2-dimensional cycles except for the triangulation of the real projective plane given in Figure 3b which is non-orientable.

5 A combinatorial condition for non-zero homology

In this section we will show that a d-dimensional cycle is the right combinatorial structure to describe the idea of non-zero d-dimensional homology over a field of characteristic 2. We will also show that orientable d-dimensional cycles produce non-zero d-homology over any field.

We begin by investigating the relationship between the combinatorial structure of a simplicial complex and its simplicial homology over $\mathbb{Z}_2$. In this field the role played by the coefficients in a 1-chain is reduced to indicating whether or not a face is present. As well since $-1 = 1$ over $\mathbb{Z}_2$ the concept of an orientation of a face is unnecessary as all orientations of a face are equivalent. This allows us to more easily examine the connection between the combinatorics of the complex and the more algebraic concepts of the 1-cycle and the 1-boundary.

The following lemma demonstrates the relationship between d-dimensional cycles and homological d-cycles over $\mathbb{Z}_2$.

Lemma 5.1 (**d-dimensional cycles are d-cycles and conversely).** *The sum of the d-faces of a d-dimensional cycle is a homological d-cycle over $\mathbb{Z}_2$ and, conversely, the d-path-connected components of the support complex of a homological d-cycle are d-dimensional cycles.*

Proof. Let Ω be a d-dimensional cycle with d-faces $F_1, \dots, F_k$. Setting $c = \sum_{i=1}^k F_i$ and applying the boundary map ∂_d over $\mathbb{Z}_2$ we have

$$\partial_d(c) = \sum_{i=1}^{k} (e_1^i + \cdots + e_{d+1}^i)$$

where $e_1^i, \dots, e_{d+1}^i$ are the $d+1$ edges of dimension $d-1$ belonging to F_i. Since the faces $F_1, \dots, F_k$ form a d-dimensional cycle each $(d-1)$-dimensional face appears in an even number of the faces $F_1, \dots, F_k$. Hence since our coefficients are in $\mathbb{Z}_2$ we have $\partial_d(c) = 0$ and so c is a d-cycle.

Let $c = F_1 + \cdots + F_k$ be a d-cycle over $\mathbb{Z}_2$. Applying the boundary map ∂_d we have

$$0 = \partial_d(F_1 + \cdots + F_k) = \sum_{i=1}^{k} (e_1^i + \cdots + e_{d+1}^i)$$

where $e_1^i, \dots, e_{d+1}^i$ are the $d+1$ edges of dimension $d-1$ belonging to F_i. If the support complex of c is not d-path-connected then we can partition this complex into n d-path-connected components $\Phi_1, \dots, \Phi_n$. Let $P_i \subseteq \{1, \dots, k\}$ be such that $F_j \in \Phi_i$ if and only if $j \in P_i$. Note that $P_1, \dots, P_n$ form a partition of $\{1, \dots, k\}$. Since $\Phi_1, \dots, \Phi_n$ have no $(d-1)$-faces in common, for each $1 \leq i \leq n$ we must have

$$\sum_{j \in P_i} (e_1^j + \cdots + e_{d+1}^j) = 0$$

and so, since our coefficients are in $\mathbb{Z}_2$, we see that each $(d-1)$-face occurring in Φ_i belongs to an even number of the d-faces in $\{F_j | j \in P_i\}$. Therefore Φ_i is a d-dimensional cycle and hence the d-path-connected components of the support complex of c all form d-dimensional cycles. ☐

Due to the close association between homological d-cycles and d-dimensional cycles over the field $\mathbb{Z}_2$ we are able to obtain a necessary and sufficient combinatorial condition for non-zero homology over any field of characteristic 2.

Theorem 5.2 (When simplicial homology vanishes in characteristic 2). *For any simplicial complex Γ and any field k of characteristic 2, $\tilde{H}_d(\Gamma; k) \neq 0$ if and only if Γ contains a d-dimensional cycle, the sum of whose d-faces is not a d-boundary.*

Proof. By application of the Universal Coefficient Theorem given in [4, Theorem 3A.3] and the property of faithful flatness we find that for any field k of characteristic 2 we have $\tilde{H}_d(\Gamma; k) \neq 0$ if and only if $\tilde{H}_d(\Gamma; \mathbb{Z}_2) \neq 0$. Therefore we need only prove the theorem in the case that $k = \mathbb{Z}_2$.

Suppose that $\tilde{H}_d(\Gamma; \mathbb{Z}_2) \neq 0$. Then Γ contains a d-cycle c that is not a d-boundary. We may assume that the support complex of c is minimal with respect to this property. In other words no strict subset of the d-faces of c has a sum that is also a d-cycle which is not a d-boundary. First we would like to show that the support complex of c is d-path connected.

Since we are in $\mathbb{Z}_2$, the d-cycle c is of the form

$$c = F_1 + \cdots + F_k$$

for some d-faces $F_1, \ldots, F_k$ of Γ. Since c is a d-cycle then applying the boundary map ∂_d we have

$$0 = \partial_d(F_1 + \cdots + F_k) = \sum_{i=1}^{k}(e_1^i + \cdots + e_{d+1}^i)$$

where $e_1^i, \ldots, e_{d+1}^i$ are the $d+1$ edges of dimension $d-1$ belonging to F_i. If the support complex of c is not d-path-connected then, without loss of generality, we can partition its set of d-faces $\{F_1, \ldots, F_k\}$ into two sets, $\{F_1, \ldots, F_\ell\}$ and $\{F_{\ell+1}, \ldots, F_k\}$ such that these two sets have no $(d-1)$-faces in common. Hence we must have

$$\sum_{i=1}^{\ell}(e_1^i + \cdots + e_{d+1}^i) = 0 \qquad \text{and} \qquad \sum_{i=\ell+1}^{k}(e_1^i + \cdots + e_{d+1}^i) = 0.$$

In other words we have

$$\partial_d(F_1 + \cdots + F_\ell) = 0 \qquad \text{and} \qquad \partial_d(F_{\ell+1} + \cdots + F_k) = 0$$

and so $F_1 + \cdots + F_\ell$ and $F_{\ell+1} + \cdots + F_k$ are both d-cycles. By our assumption of minimality these d-cycles are both d-boundaries. Hence in Γ there exist $(d+1)$-faces $G_1, \ldots, G_r$ and $H_1, \ldots, H_t$ such that

$$\partial_{d+1}\left(\sum_{i=1}^{r} G_i\right) = F_1 + \cdots + F_\ell \qquad \text{and} \qquad \partial_{d+1}\left(\sum_{i=1}^{t} H_i\right) = F_{\ell+1} + \cdots + F_k.$$

But then we have

$$\partial_{d+1}\left(\sum_{i=1}^{r} G_i + \sum_{i=1}^{t} H_i\right) = F_1 + \cdots + F_k$$

which is a contradiction since $F_1 + \cdots + F_k$ is not a d-boundary. Therefore the support complex of c must be d-path-connected. Hence, by Lemma 5.1 the support complex of c is a d-dimensional cycle. Hence Γ contains a d-dimensional cycle the sum of whose d-faces is not a d-boundary.

Conversely suppose that Γ contains a d-dimensional cycle with d-faces $F_1, \ldots, F_k$ such that $\sum_{i=1}^{k} F_i$ is not a d-boundary. By Lemma 5.1 we know that $\sum_{i=1}^{k} F_i$ is a d-cycle. It follows that $\tilde{H}_d(\Gamma; \mathbb{Z}_2) \neq 0$. $\square$

When we broaden the scope of our investigations to study simplicial homology over any field we must keep in mind examples such as the triangulation of the real projective plane given in Figure 3b. The simplicial homology of this complex changes significantly depending on the field under consideration. In particular this complex has non-zero 2-dimensional homology only over fields of characteristic 2. As we saw in Section 4 the triangulation of the real projective plane is an example of a non-orientable d-dimensional cycle. It is this notion of orientability which leads us to a sufficient condition for a simplicial complex to have non-zero homology over any field. First we see that orientable d-dimensional cycles are homological d-cycles over any field.

Lemma 5.3 (Orientable d-dimensional cycles are d-cycles). *The sum of the oriented d-faces of an orientable d-dimensional cycle is a homological d-cycle over any field k.*

Proof. Let Ω be an orientable d-dimensional cycle with d-faces $F_1, \ldots, F_k$ and let

$$c = F_1 + \cdots + F_k$$

where $F_1, \ldots, F_k$ are oriented according to the orientability of Ω. Let $e_1^i, \ldots, e_{d+1}^i$ be the $d+1$ faces of dimension $d-1$ belonging to F_i. Applying the boundary map to c we have without loss of generality

$$\partial_d(c) = \sum_{i=1}^{k} \sum_{j=1}^{d+1} (-1)^{j+1} e_j^i. \tag{3}$$

Notice that every $(d-1)$-face of Ω occurs an even number of times in (3) since Ω is a d-dimensional cycle. Furthermore, because Ω is orientable the number of times that the $(d-1)$-face appears with a positive sign is equal to the number of times that it appears with a negative sign. Hence we get $\partial_d(c) = 0$ and so c is a homological d-cycle. □

Theorem 5.4 (Orientable d-dimensional cycles give non-zero homology over all fields). *For any simplicial complex Γ and any field k, if Γ contains an orientable d-dimensional cycle the sum of whose oriented d-faces is not a d-boundary, then $\tilde{H}_d(\Gamma; k) \neq 0$.*

Proof. Let Ω be the orientable d-dimensional cycle in Γ given by our assumption and with d-faces $F_1, \ldots, F_k$. By Lemma 5.3 we know that the d-chain $c = F_1 + \cdots + F_k$ is a homological d-cycle where $F_1, \ldots, F_k$ are oriented according to the orientability of Ω. By assumption c is not a d-boundary and so $\tilde{H}_d(\Gamma; k) \neq 0$. □

The converse of this theorem does not hold as can be seen from the example of the triangulation of the real projective plane. As mentioned above this simplicial complex has non-zero 2-dimensional homology over any field of characteristic 2, but contains no orientable 2-dimensional cycles. The whole simplicial complex is, however, a non-orientable 2-dimensional cycle. Thus far we have been unable to find any counter-examples to the converse of Theorem 5.4 in the case that the field in question has characteristic 0.

References

[1] N. Biggs. *Algebraic Graph Theory*. Cambridge University Press, 2nd edition, 1993.

[2] E. Connon and S. Faridi. Chorded complexes and a necessary condition for a monomial ideal to have a linear resolution. arXiv:1209.5089, 2012.

[3] A. L. Fogelsanger. *The Generic Rigidity of Minimal Cycles*. PhD thesis, Cornell University, Ithaca, New York, May 1988.

[4] A. Hatcher. *Algebraic Topology*. Cambridge University Press, Cambridge, 2002.

[5] M. Hochster. Cohen-Macaulay rings, combinatorics, and simplicial complexes. In B. R. McDonald and R. A. Morris, editors, *Ring Theory II*, volume 26 of *Lect. Notes in Pure and Appl. Math.*, pages 171–223, New York, 1977. Dekker.

[6] G. Kalai. f-Vectors of acyclic complexes. *Discrete Math.*, 55:97–99, 1984.

[7] J. R. Munkres. *Elements of Algebraic Topology*. The Benjamin/Cummings Publishing Company, Inc., Menlo Park, California, 1984.

[8] G. A. Reisner. Cohen-Macaulay quotients of polynomial rings. *Adv. in Math.*, 21:30–49, 1976.

[9] R. P. Stanley. The upper bound conjecture and Cohen-Macaulay rings. *Stud. Appl. Math.*, 54:135–142, 1975.

[10] R. P. Stanley. A combinatorial decomposition of acyclic simplicial complexes. *Discrete Math.*, 120:175–182, 1993.

[11] D. B. West. *Introduction to Graph Theory*. Prentice-Hall, Upper Sadlle River, NJ, 1996.